



Cardinal Health™

T.E.D.™ anti-embolism stockings for continuing care

Optimal treatment for deep vein thrombosis (DVT)

Cardinal Health is proud to offer effective compression solutions for the recuperating patient. Our T.E.D.™ anti-embolism stockings are clinically proven to reduce the risk of deep vein thrombosis (DVT),¹ which can extend up to six weeks after hospitalization.^{2,3,4}

Benefits of T.E.D.™ stockings

- Clinically proven to reduce the risk of DVT¹
- Designed for the recuperating patient*
- Graduated compression pattern
- Closed-toe style for everyday use
- Available in knee length and thigh length




References: 1. Sachdeva A, Dalton M, Lees T. Graduated compression stockings for prevention of deep vein thrombosis. *Cochrane Database Syst Rev.* 2018;11:Cd001484. 2. Bratzler, et al. Development of National Performance Measures on the Prevention and Treatment of Venous Thromboembolism. *Journal of Thrombosis and Thrombolysis*, Nov. 2009. 3. White RH, Gettner S, Newman JM, Trauner KB, Romano PS (2000) Predictors of re-hospitalization for symptomatic venous thromboembolism after total hip arthroplasty. *New England Journal of Medicine* 343:1758–1764. 4. Bergqvist D, Agnelli G, Cohen AT, Eldor A, Nilsson PE, Le Moigne-Amrani A, Dietrich-Neto F, Enoxacan II Investigators (2002) Duration of prophylaxis against venous thromboembolism with enoxaparin after surgery for cancer. *New England Journal of Medicine* 346:975–980.

* Data on File


CardinalHealth
Essential to care™

Cardinal Health™ T.E.D.™ anti-embolism stockings for continuing care

Clinically-proven pressure pattern



Fitting recommendations



Knee length Measure in standing position if possible

1. Measure calf circumference at greatest portion to determine size.
2. Measure the distance from bend of knee to bottom of heel to determine length.



Thigh length Measure in standing position if possible

1. Measure upper thigh circumference at the buttock fold. When thigh circumference measures greater than 25 in., select knee length style.
2. Measure calf circumference at greatest portion to determine size.
3. Measure distance from buttock fold to base of heel to determine length.

Knee length

1. Calf circumference	2. Length	Description	Item code		
			White	Beige	Black
<12"	<16 in.	Small Regular	4278-	4265-	4434
	>16 in.	Small Long	4282	4317-	4572-
12-15 in.	<17 in.	Medium Regular	4279-	4271	4435
	>17 in.	Medium Long	4283	4323-	4573
15-17.5 in.	<18 in.	Large Regular	4280	4289	4436
	>18 in.	Large Long	4284	4335	4574
17.5-20 in.	<18 in.	X-Large Regular	4281	4296-	4437-
	>18 in.	X-Large Long	4285	4344	4575

Thigh length

1. Thigh circumference	2. Calf circumference	3. Length	Short	Regular	Long
			<29 in.	29-33 in.	>33 in.
<25 in.	Small	<12 in.	4297-	4300-	4303
	Medium	12-15 in.	4298-	4301-	4304
	Large	15-17.5 in.	4299	4302	4305-

Available in white only

T.E.D.™ stocking contraindications

Stockings may not be recommended for patients with the following:

1. Any leg condition in which stockings would interfere, such as:
 - Dermatitis (skin problems)
 - Gangrene
 - Vein ligation (recent leg surgery)
 - Recent skin graft
2. Severe vascular diseases, such as arteriosclerosis (hardening of the arteries) that cause poor circulation to the leg
3. Massive edema (swelling) of the legs, or pulmonary edema (fluid in the lungs) from congestive heart failure
4. Extreme deformity of the leg
5. For thigh length stockings, patients with a circumference greater than 25 inches at the gluteal furrow

T.E.D.™ stocking precautions

- Proper sizing and application must be assured.
- **Important:** These stockings are only to be worn with a doctor's advice and during the time period recommended.

**SMART
COMPRESSION
EDUCATES**

Join us in the fight against HA-VTE

Visit cardinalhealth.com/KnowVTE to learn more about strategies and solutions for prevention.

For more information or to order, call **800.964.5227** or visit **cardinalhealth.com/TED**

Cardinal Health
Essential to care™